



Measured Parameters


Test Name	Acronym	Units of Measure	Measurement Range	Normal Range
pH	pH	pH units	6.5 - 8.0	7.35 - 7.45 arterial
				7.32 - 7.43 venous
Carbon Dioxide, Partial Pressure	pCO ₂	mm Hg	5 - 250	35 - 48 arterial
				41 - 51 venous
		kPa	0.7 - 33.3	4.7 - 6.4 arterial
				5.4 - 6.8 venous
Oxygen, Partial Pressure	pO ₂	mm Hg	5 - 750	83 - 108 arterial
		kPa	0.7 - 100	11.1 - 14.4 arterial
Sodium	Na ⁺	mmol/L	85 - 180	138 - 146
Potassium	K ⁺	mmol/L	1.5 - 12.0	3.5 - 4.5
		mEq/L		
Ionized Calcium	Ca ⁺⁺	mmol/L	0.25 - 4.00	1.15 - 1.33
		mg/dL	1.0 - 16.0	4.6 - 5.3
		mEq/L	0.5 - 8.0	2.3 - 2.7
Chloride	Cl ⁻	mmol/L	65 - 140	98 - 107
		mEq/L		
Glucose	Glu	mmol/L	1.1 - 38.5	4.1 - 5.5
		mg/dL	20 - 700	74 - 100
		g/L	0.20 - 7.00	0.74 - 1.00
Lactate	Lac	mmol/L	0.30 - 20.00	0.56 - 1.39
		mg/dL	2.7 - 180.2	5.0 - 12.5
		g/L	0.03 - 1.80	0.05 - 0.12
Creatinine	Crea	mg/dL	0.30 - 15.00	0.51 - 1.19
		μmol/L	27 - 1326	45 - 105
Hematocrit	Hct	% PCV	10 - 75	38 - 51
		L/L	0.10 - 0.75	0.38 - 0.51

Calculated Parameters

Test Name	Acronym	Units of Measure	Measurement Range	Normal Range
Hemoglobin	cHgb	g/dL	3.3 - 25	12 - 17
		mmol/L	2.0 - 15.5	7.4 - 10.6
		g/L	33 - 250	120 - 170
Actual Bicarbonate	cHCO ₃ ⁻	mmol/L	1 - 85	21 - 28 arterial
				22 - 29 venous
		mEq/L	1 - 85	21 - 28 arterial
				22 - 29 venous
Total Carbon Dioxide	cTCO ₂	mmol/L	1 - 85	22 - 29 arterial
				23 - 30 venous
		mEq/L	1 - 85	22 - 29 arterial
				23 - 30 venous
Base Excess of Extra Cellular Fluid	BE(ecf)	mmol/L	-30 - +30	-2 - +3
		mEq/L		
Base Excess of Blood	BE(b)	mmol/L	-30 - +30	-2 - +3
		mEq/L		
Oxygen Saturation	cSO ₂	%	0 - 100	94 - 98
Estimated Glomerular Filtration Rate	eGFR	mL/min/1.73m ²	2 - 60 or >60	**
Estimated Glomerular Filtration Rate if African American	eGFR-a	mL/min/1.73m ²	2 - 60 or >60	**
** Institutions should establish and set their own normal range values				
Anion Gap	AGap	mmol/L	-14 - +95	7 - 16
		mEq/L		
Anion Gap, K ⁺	AGapK	mmol/L	-10 - +99	10 - 20
		mEq/L		

Alere For more information, please visit alere-epoc.com

© 2016 Alere. All rights reserved. The Alere Logo, Alere, epoc and Knowing now matters are trademarks of the Alere group of companies. 10002171-03 06/16
All other trademarks referenced are trademarks of their respective owners.



epoc® Blood Analysis System

The Right Result at the Right Time

Gain immediate access to critical test results.

The **epoc® System** is a handheld, wireless solution that provides blood gas, electrolyte and metabolite results at the patient's bedside in approximately 30 seconds after sample introduction.



Test Card

- Room temperature storage
- Barcoded with lot and expiration for error-free test panel recognition
- 92 µL (about a 1/10th of a CC) sample

11 critical tests on a single card

pH	pO ₂	K ⁺	Cl ⁻	Lac	Hct
pCO ₂	Na ⁺	Ca ⁺⁺	Glu	Crea	

Calculated values

AGap	cHCO ₃ ⁻	BE(ecf)	cSO ₂	eGFR
AGapK	cTCO ₂	BE(b)	cHgb	eGFR-a

epoc Host² Mobile Computer

- Runs the customizable epoc Host Software and is the caregiver's interface to the system
- Customizable reference and critical ranges
- Electronic documentation of test results, patient information and respiratory parameters

epoc Reader

- Auto-calibrates the epoc Test Card prior to sample introduction
- Communicates bi-directionally with the epoc Host via Bluetooth®
- Fully portable; AC/rechargeable battery

epoc Enterprise Data Manager (EDM) Software

- Provides access through any computer on the hospital network
- Seamlessly connects to hospital LIS/HIS via industry standard HL7® interface
- Offers the ability to manage the entire epoc System, interface and users

Streamline the patient testing process

By moving patient testing to the bedside, caregivers, patients and hospital administration all benefit from a more streamlined process with improved turn around times and operational efficiencies.

epoc® System 7 Steps ~3 Minutes

Portable / Near Care Testing 14 Steps 15-20 Minutes

Central Lab Testing 18 Steps 20 Minutes-2 hours



Wireless communication delivers real-time results and reporting

With the epoc System, easily transmit patient results from the bedside to the electronic medical record (LIS/HIS) in seconds. Reduce transcription errors or delays when downloading results at remote locations.

